

# The non-absoluteness of model existence in uncountable cardinals for $L_{\omega_1, \omega}$

Sy-David Friedman (KGRC, Vienna)\*  
Tapani Hyttinen (University of Helsinki) †  
Martin Koerwien (CRM, Barcelona) ‡

May 3, 2012

## Abstract

For sentences  $\phi$  of  $L_{\omega_1, \omega}$ , we investigate the question of absoluteness of  $\phi$  having models in uncountable cardinalities. We first observe that having a model in  $\aleph_1$  is an absolute property, but having a model in  $\aleph_2$  is not as it may depend on the validity of the Continuum Hypothesis. We then consider the GCH context and provide sentences for any  $\alpha \in \omega_1 \setminus \{0, 1, \omega\}$  for which the existence of a model in  $\aleph_\alpha$  is non-absolute (relative to large cardinal hypotheses). Finally, we present a *complete* sentence for which model existence in  $\aleph_3$  is non-absolute.

Keywords: Infinitary Logic, Absoluteness, Abstract Elementary Classes

2010 MSC: 03C48, 03C75

Throughout, we assume  $\phi$  is an  $L_{\omega_1, \omega}$  sentence which has infinite models. By the downward Löwenheim-Skolem-Theorem,  $\phi$  must have a countable model, so the property “having a countable model” is an absolute property of such sentences in the sense that its validity does not depend on the properties of the set-theoretic universe we work in. More precisely, if  $V \subseteq W$  are transitive models of ZFC with the same ordinals and  $\phi \in V$ ,  $V \models$  “ $\phi$  is an  $L_{\omega_1, \omega}$ -sentence” (with a natural set-theoretic coding of such sentences), then  $V \models$  “ $\phi$  has a countable model” if and only if  $W \models$  “ $\phi$  has a countable model”. The purpose of this paper is to investigate the question of how far we can replace “countable” by higher cardinalities.

---

\*Supported by the John Templeton Foundation under grant number 13152, *The Myriad Aspects of Infinity* and by the Austrian Science Fund (FWF) through Project Number P 22430-N13.

†Partially supported by the Academy of Finland under grant number 1123110.

‡Supported by the John Templeton Foundation under grant number 13152, *The Myriad Aspects of Infinity*.

A main tool for absoluteness considerations is Shoenfield's absoluteness Theorem (Theorem 25.20 in [9]). It states that any property expressed by either a  $\Sigma_2^1$  or a  $\Pi_2^1$  formula is absolute between transitive models of ZFC with the same ordinals. As John Baldwin observed in [1], it follows from results of [7] that the property of  $\phi$  having arbitrarily large models is absolute (it can be expressed in form of the existence of an infinite indiscernible sequence, which by Shoenfield is absolute). Since the Hanf number of the logic  $L_{\omega_1, \omega}$  equals  $\beth_{\omega_1}$ , it follows that the existence of models in cardinalities above that number is absolute. Therefore the context we are interested in is where  $\phi$  (absolutely) does not have a model of size  $\beth_{\omega_1}$ .

## 1 The case $\aleph_1$

For *complete* sentences  $\phi$  (meaning that any model of  $\phi$  satisfies the same  $L_{\omega_1, \omega}$  sentences), having a model in  $\aleph_1$  is an absolute notion. We have the following characterization (which appears also in [1] as well as [5]) of  $\phi$  having a model of size  $\aleph_1$  (which is a  $\Sigma_1^1$  property and therefore absolute by Shoenfield's absoluteness Theorem):

- (\*) There exist two countable models  $M, N$  of  $\phi$  such that  $M$  is a proper elementary (in the fragment of  $\phi$ ) substructure of  $N$ .

To see that this is a characterization, note first that if  $\phi$  has an uncountable model, (\*) holds by Löwenheim-Skolem. For the converse, we use the completeness of  $\phi$  which implies that any two countable models of  $\phi$  are isomorphic (by Scott's isomorphism Theorem, since  $\phi$  is complete and thus characterizes its countable models up to isomorphism). Then, as  $N \cong M$ , we can find a proper countable  $L_{\omega_1, \omega}$ -elementary extension of  $N$  as well and continue this procedure  $\omega_1$  many times (taking unions at limit stages). The union of this elementary chain will then be a model of  $\phi$  of size  $\aleph_1$ .

If the sentence is not complete, criterion (\*) does not obviously imply the existence of an uncountable model. By a theorem of Gregory's (see [6]), it can be seen that it actually does. We will however provide a different criterion ((\*\*) below) for which we have a relatively basic proof (essentially only using the omitting types theorem for  $L_{\omega_1, \omega}$ ) that it is equivalent to  $\phi$  having an uncountable model. We have thus that for any (even incomplete)  $L_{\omega_1, \omega}$ -sentence, model existence in  $\aleph_1$  is absolute<sup>1</sup>.

In the following, we consider the sentence  $\phi$  as a set-theoretic object using standard coding of formulas of  $L_{\omega_1, \omega}$ .  $\phi$  can thus be regarded as a hereditarily countable set.

The following property which (again by Shoenfield) is absolute, characterizes  $\phi$  having a model of size  $\aleph_1$ :

---

<sup>1</sup>This has also been observed recently by Paul Larson. His argument uses iterated generic ultrapowers. Rami Grossberg points out, he knew of this fact already in the 1980's but did not publish it, and that others like Shelah, Barwise and Keisler most likely knew of it even earlier.

(\*\*) There is a countable transitive model  $U$  of  $\text{ZFC}^-$  (ZFC without the power set axiom) containing  $\phi$  with  $U \models$  “ $\omega_1$  exists,  $\phi$  is hereditarily countable, and there is a model of  $\phi$  with universe  $\omega_1$ ”.

First, suppose  $\phi$  has a model  $M$  of size  $\aleph_1$ , say one with universe  $\omega_1$ . As both  $\phi$  and  $M$  are elements of  $H_{\omega_2}$  (the collection of sets hereditarily of size at most  $\aleph_1$ ), we have  $H_{\omega_2} \models \text{ZFC}^- +$  “there is a model of  $\phi$  with universe  $\omega_1$ ”. Now it suffices to take a countable (first order) elementary substructure  $U \prec H_{\omega_2}$  containing  $\phi$ , and  $U$  will have the properties of (\*\*).

Conversely, assuming (\*\*) holds for some countable  $U$ , we can take an elementary extension  $U'$  of  $U$  where all (in the sense of  $U$ ) hereditarily countable sets are unchanged and all (in  $U$ ) uncountable ones become sets of size  $\aleph_1$ . This can be achieved using Corollary A of Theorem 36 in [11], noting that it holds for models of  $\text{ZFC}^-$  (instead of full ZFC as the Corollary originally assumes), as the powerset axiom is not used for it. In particular this is true for the  $\omega_1$  of  $U'$  on which we know a model  $M$  of  $\phi$  lives (note that  $U' \models (M \models \phi)$  implies that  $M \models \phi$  in the real universe; to see this, use that  $U'$  contains the fragment of  $\phi$  and satisfaction for formulas in this fragment is absolute between  $U'$  and the real universe”). So we get a model of  $\phi$  of size  $\aleph_1$ .

There is another absolute criteria characterizing  $\phi$  having an uncountable model, but it requires going beyond the logic  $L_{\omega_1, \omega}$ . Let us consider the extension  $L_{\omega_1, \omega}(Q)$  of  $L_{\omega_1, \omega}$  obtained by adding an extra quantifier  $Q$  with the semantics “there exist uncountably many”. As is shown in [2],  $L_{\omega_1, \omega}(Q)$  admits a completeness theorem which actually has a very natural (absolute) deduction calculus. Now the statement

(\*\*\*) There is a proof of  $\neg Qx(x = x)$  starting from  $\phi$

characterizes  $\phi$  having only countable models. Thus the negation of (\*\*\*) is an (absolute) property characterizing  $\phi$  having an uncountable model. Note that this argument shows that model existence in  $\aleph_1$  is absolute even for  $L_{\omega_1, \omega}(Q)$  sentences.

## 2 Going beyond $\aleph_1$

It is not generally true that the existence of a model of size  $\aleph_2$  is an absolute property.

A very simple way to see this is to take any sentence  $\phi$  that has models exactly up to size continuum. We easily find even *complete* sentences with this property. Then clearly,  $\phi$  has a model of size  $\aleph_2$  if and only if the continuum hypothesis fails.

More generally, such a sentence has a model of size  $\aleph_\alpha$  if and only if  $2^{\aleph_0} \geq \aleph_\alpha$ . So for any  $\alpha > 1$ , the existence of a model of size  $\aleph_\alpha$  is non-absolute.

There are many examples of complete  $L_{\omega_1, \omega}$ -sentences in the literature having models exactly up to size continuum, but they are mostly more complicated than necessary

for our purposes, because their authors have been interested in additional properties. Therefore we provide here a very simple such example which uses the idea of coding full binary trees. This same idea has been used in Malitz's examples showing that the Hanf number for *complete*  $L_{\omega_1, \omega}$ -sentences equals  $\beth_{\omega_1}$  (see [12]).

Let the language  $L$  consist of countably many binary relation symbols  $E_n$  ( $n < \omega$ ), and let  $\sigma \in L_{\omega_1, \omega}$  be the conjunction of

- All  $E_n$  are equivalence relations such that  $E_0$  has two classes and each  $E_n$ -class is the union of exactly two  $E_{n+1}$ -classes.
- $\forall x, y ((\bigwedge_{n < \omega} E_n(x, y)) \rightarrow x = y)$

It is an easy back-and-forth argument to show that any two countable models of  $\sigma$  are isomorphic, so  $\sigma$  is complete. Every model represents a set of branches through a full binary tree, so there cannot be models greater than the continuum. On the other hand, the Cantor space  $2^\omega$  together with the relations “ $E_n(x, y)$  if and only if  $x$  and  $y$  coincide on the  $n + 1$  first components” is a model of  $\sigma$  of size continuum.

### 3 Going beyond $\aleph_1$ under the assumption of GCH

As we have seen, playing with the cardinal exponential function provides trivial examples for the non-absoluteness of the existence of models of cardinality greater than  $\aleph_1$ . A next natural question is if this is the only non-absoluteness phenomenon there is. That is, under the additional assumption of GCH, does the existence of models in cardinalities greater than  $\aleph_1$  become an absolute notion? We will provide different incomplete sentences and later on even a complete one that show the answer is negative.

#### 3.1 A reminder about two-cardinal properties

As we will see later, there is an interesting connection between classical first-order two-cardinal properties and model existence for  $L_{\omega_1, \omega}$ -sentences. We recall the following definition:

**Definition 1.** *Let  $T$  be a first-order theory in a signature containing a unary predicate  $P$ . Given two infinite cardinals  $\kappa \geq \lambda$ , we say that  $T$  admits  $(\kappa, \lambda)$  if there is a model of  $T$  of size  $\kappa$  such that  $P^M = \{a \in M \mid M \models P(a)\}$  has cardinality  $\lambda$ .*

As is already exposed in Chang-Keisler's classical textbook [3] in chapter 7.2, admitting certain pairs  $(\kappa, \lambda)$  is a non-absolute property for certain theories. There, examples are given where admitting  $(\kappa^+, \kappa)$  is equivalent to the existence of a special  $\kappa^+$ -Aronszajn tree or where admitting  $(\kappa^{++}, \kappa)$  is equivalent to the existence of a  $\kappa^+$  Kurepa tree (or equivalently a  $\kappa^+$  Kurepa family).

### 3.2 Some set theory

We now recall the two classical concepts of Kurepa families and special Aronszajn trees. The first-order examples in [3] showing non-absoluteness of the existence of certain two-cardinal models and our later exposed examples of  $L_{\omega_1, \omega}$ -sentences showing non-absoluteness of model existence in certain cardinalities code those objects in their models. The coding is such that the existence of a certain two-cardinal model or the existence of a model in a certain cardinality is equivalent to the existence of such an object (which is independent from ZFC+GCH as we will see in the following).

**Definition 2.** *Let  $\kappa$  be any infinite cardinal. A  $\kappa^+$  Kurepa family is a family  $\mathcal{F}$  of subsets of some set  $A$  with  $|A| = \kappa^+$ , such that  $|\mathcal{F}| > \kappa^+$  and for any subset  $B \subset A$  with  $|B| = \kappa$ ,  $|\{X \cap B \mid X \in \mathcal{F}\}| \leq \kappa$ .*

*Let  $\text{KH}_{\kappa^+}$  be the statement that there exists a  $\kappa^+$  Kurepa family.*

It is folklore that the existence of Kurepa families in different  $\aleph_\alpha$  ( $\alpha < \omega_1$ ) is independent from one another. We will now describe the formal arguments for the cases we need (essentially the same arguments would work more generally for “switching on and off” independently the existence of Kurepa families in different  $\aleph_\alpha$ ). In the constructible universe,  $\text{KH}_{\kappa^+}$  is true for all cardinals  $\kappa$  (this follows from the fact that  $\diamond^+$  holds at successor cardinals in  $L$ , see [10]). On the other hand we have:

**Theorem 3.** *The consistency of “ZFC+there are uncountably many inaccessible cardinals” implies the consistency of “ZFC+GCH+ $\forall \alpha < \omega_1 \neg \text{KH}_{\aleph_{\alpha+1}}$ ”*

*Proof.* This is a slight generalisation of Silver’s argument that if  $\kappa$  is inaccessible then after forcing with  $\text{Coll}(\omega_1, < \kappa)$ , the forcing to convert  $\kappa$  into  $\aleph_2$  with countable conditions,  $\text{KH}_{\aleph_1}$  fails (see [9], Theorem 27.9).

Assume GCH, let  $\kappa_0$  be  $\aleph_1$  and define  $(\kappa_\beta)_{0 < \beta < \omega_1}$  inductively: set  $\kappa_{\beta+1}$  the least inaccessible cardinal greater than  $\kappa_\beta$  and for  $\beta < \omega_1$  a limit ordinal set  $\kappa_\beta = \sup\{\kappa_\gamma \mid \gamma < \beta\}^+$ . Let  $P$  be the fully supported product of the forcings  $\text{Coll}(\kappa_\beta, < \kappa_{\beta+1})$  for  $\beta < \omega_1$ . Then in the extension,  $\kappa_\beta$  equals  $\aleph_{\beta+1}$ , while the GCH still holds. We claim that  $\text{KH}_{\kappa_\beta}$  fails for each  $\beta < \omega_1$ .

Indeed, the forcing  $P$  can be factored as  $P(< \beta) \times P(\geq \beta)$  where  $P(< \beta)$  refers only to the collapses  $\text{Coll}(\kappa_\gamma, < \kappa_{\gamma+1})$  for  $\gamma < \beta$  and  $P(\geq \beta)$  refers only to the collapses  $\text{Coll}(\kappa_\gamma, < \kappa_{\gamma+1})$  for  $\gamma \geq \beta$ . Similarly,  $V[G]$  factors as  $V[G(< \beta)][G(\geq \beta)]$ . In the model  $V[G(< \beta)]$ ,  $\kappa_{\beta+1}$  is still inaccessible, so we can apply Silver’s argument to conclude that  $\text{KH}_{\kappa_\beta}$  fails in  $V[G(< \beta)][G(\geq \beta)] = V[G]$ , using the closure of the forcing  $P(\geq \beta)$  under sequences of length less than  $\kappa_\beta$ .  $\square$

**Definition 4.** *A tree is a partially ordered set  $(T, <)$  such that for any element  $t \in T$ , the set  $\{x \mid x < t\}$  is well ordered by  $<$ . The rank  $\text{rk}(t)$  of  $t$  is the order type of  $\{x \mid x < t\}$ . For any ordinal  $\alpha$ , let  $T_\alpha = \{t \in T \mid \text{rk}(t) = \alpha\}$ .*

For any cardinal  $\kappa$ , a  $\kappa^+$ -tree is a tree  $T$  such that  $T_{\kappa^+} = \emptyset$  and for all  $\alpha < \kappa^+$ ,  $0 < |T_\alpha| < \kappa^+$ .  $T$  is normal, if

- $|T_0| = 1$
- every element has at least two immediate successors
- for any  $t \in T$  and  $\alpha$  with  $\text{rk}(t) < \alpha < \kappa^+$ , there is some  $t' > t$  with  $\text{rk}(t') = \alpha$ .

A normal  $\kappa^+$ -tree  $T$  is a special  $\kappa^+$ -Aronszajn tree, if there is some set  $A$  of size  $\kappa$  and a function  $f : T \rightarrow A$  such that for all  $t, t' \in T$ ,  $t < t'$  implies  $f(t) \neq f(t')$ .

It is a consequence of GCH that special  $\kappa$ -Aronszajn trees exist for all successor cardinals  $\kappa$  that are not successors of singular cardinals (see [13]). Moreover, in the constructible universe, special Aronszajn trees exist even in successors of singular cardinals (this is a consequence of  $\square_\kappa$ , see [10]).

On the other hand, the consistency of “ZFC+ $\exists\kappa(\kappa$  supercompact)” implies the consistency of “ZFC+GCH+there are no special  $\aleph_\alpha$ -Aronszajn trees for all countable limit successors  $\alpha$ ”:

We start with a model of GCH with a supercompact cardinal  $\kappa$  and force with  $\text{Coll}(\omega_1, < \kappa)$ . As is argued in [4], this forcing preserves a stationary reflection property sufficient to ensure that Weak Square fails at  $\aleph_\lambda$  for  $\lambda$  a limit ordinal of countable cofinality. By a result of Jensen in [10], Weak Square at a cardinal  $\kappa$  is equivalent to the existence of a special Aronszajn tree on  $\kappa^+$ .

### 3.3 Connecting first-order two-cardinal properties with $L_{\omega_1, \omega}$ -model existence

We will describe how a first-order theory  $T$  can be turned into an  $L_{\omega_1, \omega}$ -sentence  $\sigma$  in such a way that  $T$  admitting certain  $(\kappa, \lambda)$  is equivalent to the existence of a model of  $\sigma$  of size  $\kappa$ .

We start with the definition of an  $L_{\omega_1, \omega}$ -sentence  $\sigma_0^\alpha$  characterizing  $\aleph_\alpha$  (for  $\alpha < \omega_1$ ), which means that it (absolutely) has a model of size  $\aleph_\alpha$ , but no bigger model. We wish to point out that the idea we use here of characterizing cardinals using  $\kappa$ -like orderings for various  $\kappa$  is not new. Also, there exist other ways of characterizing cardinals in the literature, most notably Hjorth’s examples presented in [8] that are even *complete* sentences.

Let  $L_0^\alpha = \{Q_\beta, a_n, <, F\}_{\beta \leq \alpha; n < \omega}$ , where the  $Q_\beta$  are unary predicates, the  $a_n$  are constant symbols,  $<$  is a binary and  $F$  a ternary relation symbol.

Let  $\sigma_0^\alpha \in (L_0^\alpha)_{\omega_1, \omega}$  be the conjunction of the following sentences:

- The universe is the union of all  $Q_\beta$

- $Q_0 = \{a_n | n < \omega\}$  where all  $a_n$  designate distinct elements.
- For any  $\beta < \alpha$ ,  $Q_{\beta+1}$  is disjoint from any  $Q_\gamma$  for all  $\gamma \leq \beta$ .
- For any limit ordinal  $\beta \leq \alpha$ ,  $Q_\beta = \bigcup_{\gamma < \beta} Q_\gamma$
- $<$  linearly orders  $Q_{\beta+1}$  for every  $\beta < \alpha$  and  $x < y$  implies that for some  $\beta < \alpha$ , both  $x$  and  $y$  belong to  $Q_{\beta+1}$ .
- $F(a, b, c)$  implies that for some  $\beta < \alpha$ ,  $a \in Q_{\beta+1}$ ,  $b < a$  and  $c \in Q_\beta$ .
- For every  $\beta < \alpha$  and every  $a \in Q_{\beta+1}$ ,  $F(a, \cdot, \cdot)$  defines a total injective function from  $\{x | x < a\}$  into  $Q_\beta$ .

Note that for  $\beta$  a limit ordinal or zero,  $Q_\beta$  is not ordered by  $<$  and if  $\alpha = 0$ , both  $<$  and  $F$  are empty relations.

Clearly, if  $M \models \sigma_0^\alpha$ , then in  $M$  the ordering of  $Q_{\beta+1}$  must be  $|Q_\beta|$ -like (i.e. any proper initial segment has cardinality at most  $|Q_\beta|$ ). This implies that  $|Q_{\beta+1}|$  is at most  $|Q_\beta|^+$  and since  $Q_0$  is countable by definition, we see inductively that the cardinality of each  $Q_\beta$  is bounded by  $\aleph_\beta$ . Also, there clearly exist models such that  $|Q_\beta| = \aleph_\beta$  for all  $\beta \leq \alpha$ .

Now suppose we have a first-order theory  $T$  in a language containing a unary predicate  $P$ . For  $\beta < \alpha < \omega_1$ , we define the  $L_{\omega_1, \omega}$ -sentence  $\sigma_T^{\alpha, \beta}$  as the conjunction of

- $T$
- $\sigma_0^\alpha$
- $P = Q_\beta$

**Proposition 5.** *Let  $\beta < \omega_1$  and  $0 < n < \omega$ .  $T$  admits  $(\aleph_{\beta+n}, \aleph_\beta)$  if and only if  $\sigma_T^{\beta+n, \beta}$  has a model of cardinality  $\aleph_{\beta+n}$ .*

*Proof.* If  $M \models \sigma_T^{\beta+n, \beta}$  has cardinality  $\aleph_{\beta+n}$ , we must have  $|Q_\beta| = \aleph_\beta$  in that model (here we use that  $n$  is finite!). Now the reduct of  $M$  to the language of  $T$  is a model of size  $\aleph_{\beta+n}$  where  $P$  has size  $\aleph_\beta$ .

Conversely, given a model of  $T$  of size  $\aleph_{\beta+n}$  where  $P$  has size  $\aleph_\beta$ , it is easy to expand this model to be a model of  $\sigma_T^{\beta+n, \beta}$ .  $\square$

Note that this Proposition becomes false if  $n$  is allowed to be infinite.

### 3.4 Examples of incomplete sentences: successor cardinals

We quote Chang-Keisler's results 7.2.11 and 7.2.13 from [3] (adapting the notation slightly):

- There is a sentence  $\phi_1$  in a finite language  $L$  such that for all infinite cardinals  $\lambda$ ,  $\phi_1$  admits  $(\lambda^+, \lambda)$  if and only if there exists a special  $\lambda^+$ -Aronszajn tree.
- There is a sentence  $\phi_2$  in a suitable language such that for all infinite cardinals  $\lambda$ ,  $\phi_2$  admits  $(\lambda^{++}, \lambda)$  if and only if a  $\lambda^+$  Kurepa family exists.

From the preceding section we get thus infinitary sentences  $\sigma_{\phi_1}^{\alpha+1, \alpha}$  and  $\sigma_{\phi_2}^{\alpha+2, \alpha}$  such that

- $\sigma_{\phi_1}^{\alpha+1, \alpha}$  has a model of cardinality  $\aleph_{\alpha+1}$  if and only if a special  $\aleph_{\alpha+1}$ -Aronszajn tree exists.
- $\sigma_{\phi_2}^{\alpha+2, \alpha}$  has a model of cardinality  $\aleph_{\alpha+2}$  if and only if an  $\aleph_{\alpha+1}$  Kurepa family exists.

Now recalling the set-theoretic facts from section 3.2, we get the following results:

**Theorem 6.** *Let  $\alpha < \omega_1$  be a limit ordinal. Assuming ZFC, GCH and the existence of a supercompact cardinal, model-existence in  $\aleph_{\alpha+1}$  is non-absolute for  $L_{\omega_1, \omega}$ -sentences.*

**Theorem 7.** *Let  $\alpha < \omega_1$ . Assuming ZFC, GCH and the existence of uncountably many inaccessible cardinals, model-existence in  $\aleph_{\alpha+2}$  is non-absolute for  $L_{\omega_1, \omega}$ -sentences.*

At this point, we have covered all cases of successor cardinals  $\aleph_\alpha$  for  $1 < \alpha < \aleph_{\omega_1}$ .

### 3.5 Examples of incomplete sentences: limit cardinals

We would also like to find examples of (incomplete) sentences where model existence in  $\aleph_\alpha$  is non-absolute modulo ZFC+GCH for countable limit ordinals  $\alpha$ . With a slight variation of our examples involving special Aronszajn trees, we can deal with limits that are greater than  $\omega$ .

Since the construction is rather straightforward, we will only give an informal description of it.

The sentence  $\phi_1$  used to prove Theorem 6 which is given explicitly in [3] involves essentially a binary relation  $T$  coding a tree and a unary predicate  $U$  and has the property that whenever  $M \models \phi_1$  and  $|M| = |U^M|^+$ , then  $T$  has a subtree which is a special  $|M|$ -Aronszajn-tree.

Now, fixing some  $\alpha < \omega_1$  greater than  $\omega$ , we start with the sentence  $\sigma_0^\alpha$  (see section 3.3) and for all  $\beta < \alpha$ , we add the sentence  $\phi_1$  relativised to  $\bigcup_{\gamma \leq \beta+1} Q_\gamma$  (i.e. the set

$\bigcup_{\gamma \leq \beta+1} Q_\gamma$  with the induced structure in the language of  $\phi_1$  is a model of  $\phi_1$ ) with  $Q_\beta$  taking the role of  $U$ . I.e. we are coding special Aronszajn trees at *every* level  $Q_{\beta+1}$  where  $|Q_{\beta+1}| = |Q_\beta|^+$ .

The result is a sentence  $\sigma_1^\alpha$  for which (assuming consistency of supercompact cardinals) the existence of a model of size  $\aleph_\alpha$  is non-absolute modulo ZFC+GCH. The reason is

that if no special  $\aleph_{\omega+1}$ -Aronszajn tree exists, the maximum cardinality of a model of  $\sigma_1^\alpha$  is  $\aleph_\omega$  since whenever for some  $\gamma < \alpha$ ,  $|Q_{\gamma+1}| = |Q_\gamma|^+ = \aleph_{\omega+1}$ , a special  $\aleph_{\omega+1}$ -Aronszajn tree will be coded in the model. Note that in any case,  $\sigma_1^\alpha$  will have models of size  $\aleph_\omega$  since GCH implies the existence of special  $\alpha_n$ -Aronszajn trees for all finite  $n > 0$ . Therefore these examples do not show non-absoluteness of model existence in  $\aleph_\omega$ .

## 4 A complete sentence

Both the first-order examples from [3] and our  $L_{\omega_1, \omega}$ -examples from the preceding section are highly incomplete (i.e. many first-order or  $L_{\omega_1, \omega}$ -statements are undecided) and it seems a very non-trivial task to turn them into complete theories while conserving the properties that matter to us.

We will now introduce a method of completing incomplete  $L_{\omega_1, \omega}$ -sentences that has the benefits of providing fairly explicit axiomatizations as well as some means of constructing models of the resulting complete sentence with certain properties. This method will then be applied to an incomplete sentence coding  $\aleph_2$  Kurepa trees (similar to the examples from the preceding section).

**Definition 8.** *Let  $\sigma \in L_{\omega_1, \omega}$ .*

- A  $\sigma$ -chain is a family  $(M_\alpha)_{\alpha < \lambda}$  of models of  $\sigma$  such that whenever  $\alpha < \beta < \lambda$ , we have  $M_\alpha \subset M_\beta$ .
- $\sigma$  is preserved under chains if for any  $\sigma$ -chain  $(M_\alpha)_{\alpha < \lambda}$ ,  $M = \bigcup_{\alpha < \lambda} M_\alpha$  is a model of  $\sigma$ .

As in the classical first-order case, it is still true that any  $\Pi_2$ -sentence is preserved under chains, i.e. any sentence of the form  $\forall \bar{x} \exists \bar{y} \psi(\bar{x}, \bar{y})$ , where  $\psi$  is quantifier-free (but possibly infinitary). We have to be a little careful with the definition of  $\Pi_2$  as for example infinite disjunctions of universal formulas might not be preserved under chains. A simple example is given by the sentence

$$\sigma = \bigvee_{S \subset \omega \text{ finite}} \forall x (U(x) \leftrightarrow \bigvee_{i \in S} x = a_i)$$

in the language of countably many constants  $a_i$  and a unary predicate  $U$ . This sentence expresses that  $U$  is finite.

**Definition 9.** *Let  $\sigma \in L_{\omega_1, \omega}$ .*

- Set  $S_{\text{qf}}(\sigma) = \{\text{tp}_{\text{qf}}(\bar{a}) \mid \exists M \models \sigma (\bar{a} \in M)\}$  (where  $\text{tp}_{\text{qf}}(\bar{a})$  is the quantifier-free type of  $\bar{a}$ ).
- $\sigma$  is qf-small if  $S_{\text{qf}}(\sigma)$  is countable.

Note that by the downward Löwenheim-Skolem-Theorem, we can define  $S_{\text{qf}}(\sigma)$  by referring only to countable models of  $\sigma$ .

**Definition 10.** *Suppose  $\sigma$  is qf-small.*

- For any pair  $p(\bar{x}), q(\bar{x}\bar{y}) \in S_{\text{qf}}(\sigma)$  define the sentence  $\sigma_{p,q} = \forall \bar{x}(p(\bar{x}) \rightarrow \exists \bar{y} q(\bar{x}\bar{y}))$ .
- Set  $\sigma^* = \sigma \wedge \bigwedge_{p,q \in S_{\text{qf}}(\sigma); p \subset q} \sigma_{p,q}$

If  $\sigma$  is preserved under chains, then  $\sigma^*$  is as well. However, there are consistent  $\sigma$  for which  $\sigma^*$  is inconsistent. An example would be the sentence  $\sigma = \forall a, b, c, d(R(a, b) \wedge R(c, d) \rightarrow a = c \wedge b = d)$  which expresses that exactly two points are  $R$ -related.

**Proposition 11.** *For any  $\sigma$ , if  $\sigma^*$  is consistent, it is complete.*

*Proof.* We show  $\aleph_0$ -categoricity. Let  $M, N \models \sigma^*$  be countable and suppose  $f$  is a finite partial isomorphism mapping a tuple  $\bar{a} \in M$  to a tuple  $\bar{b} \in N$ . Now let  $c \in M$  be any point and set  $p = \text{tp}_{\text{qf}}(\bar{a}) (= \text{tp}_{\text{qf}}(\bar{b}))$  and  $q = \text{tp}_{\text{qf}}(\bar{a}c)$ . Since  $N \models \sigma_{p,q}$ , we find a  $d \in N$  with  $\bar{b}d \models q$ , so we can extend  $f$  by mapping  $c$  to  $d$ . Now after enumerating both  $M$  and  $N$  we can construct a total isomorphism as the union of finite partial isomorphisms by adding every point of  $M$  to the domain and every point of  $N$  to the range eventually.  $\square$

**Definition 12.** *A sentence  $\sigma \in L_{\omega_1, \omega}$  has the extension property for countable models (EPC), if for any countable  $M \models \sigma$  and  $p(\bar{x}) \subset q(\bar{x}\bar{y})$  in  $S_{\text{qf}}(\sigma)$ , whenever some  $\bar{a} \in M$  realizes  $p$ , there is a countable  $N \models \sigma$  with  $M \subset N$  containing some  $\bar{b}$  with  $\bar{a}\bar{b} \models q$ .*

**Theorem 13.** *Suppose  $\sigma \in L_{\omega_1, \omega}$  is preserved under chains, is qf-small and has the EPC. Then*

1.  $\sigma^*$  is consistent.
2. any countable model of  $\sigma$  has an extension that is a model of  $\sigma^*$ .
3.  $\sigma^*$  is the only completion of  $\sigma$  with property 2 that is still preserved under chains.

*Proof.* Let  $M \models \sigma$  be countable. Enumerate all possible pairs  $(\bar{a}, q)$  where  $\bar{a} \in M$  and  $\text{tp}_{\text{qf}}(\bar{a}) \subset q \in S_{\text{qf}}(\sigma)$  as  $((\bar{a}_n, q_n))_{n < \omega}$ . Construct a  $\subset$ -chain  $(M_n)_{n < \omega}$  of models of  $\sigma$  such that in  $M_n$  we add a tuple  $\bar{b}_n$  with the property that  $\bar{a}_n \bar{b}_n \models q_n$ . Let  $M^1 = \bigcup_{n < \omega} M_n$ . Do

the same procedure for  $M^1$  in place of  $M$  to get some  $M^2$ . Repeat this  $\omega$  many more times and set  $N = \bigcup_{k < \omega} M^k$ . Since  $\sigma$  is preserved under chains we still have  $N \models \sigma$ , and we just added all necessary witnesses in the chains to satisfy all  $\sigma_{p,q}$  as well, so we have constructed a model of  $\sigma^*$  that contains the model  $M$  we started with.

The uniqueness of  $\sigma^*$  follows from the fact that if some  $\tau$  has the same properties, including being preserved under chains, we can form a  $\subset$ -chain  $(M_n)_{n < \omega}$  with  $M_{2n} \models \sigma^*$  and  $M_{2n+1} \models \tau$  for all  $n$ . Then by preservation under chains, the union must be a model of both  $\sigma^*$  and  $\tau$  and we conclude by completeness of both sentences.  $\square$

Now we turn to the definition of an incomplete sentence coding  $\aleph_2$  Kurepa families, which we will then complete by the described technique.

Our language will be  $\mathcal{L} = \{S, L, U, V, E_n, <, R, F, G, H\}_{n < \omega}$ , where  $S$  and  $L$  are unary predicates, all  $E_n$  as well as  $U, V, <, R$  are binary relations and  $F, G$  and  $H$  are ternary relations.

Before we give the formal definition of our sentence, we describe informally what a model of it looks like:

- $(L, <)$  is a linear order.
- The elements of  $S$  code subsets of  $L$  via the relation  $R$  such that any two of them coincide on an initial segment of  $L$  with a maximum element and are disjoint above that initial segment.
- $F$  defines a binary function mapping two elements of  $S$  to the point of  $L$  where they become disjoint.
- For every  $a \in L$ ,  $U$  and  $V$  define sets  $U_a = \{x | U(a, x)\}$ ,  $V_a = \{x | V(a, x)\}$  and all those sets are pairwise disjoint.
- The  $E_n$  are such that every set  $U_a$  and  $V_a$  with the restrictions of the  $E_n$  satisfies the theory of binary splitting equivalence relations, given in section 2. In particular, all these sets have size at most  $2^{\aleph_0} = \aleph_1$ .
- $G$  codes bijections between every initial segment  $\{x | x < a\}$  and the set  $U_a$ . This makes  $(L, <)$   $\aleph_2$ -like.
- $H$  codes intersections of sets coded by elements of  $S$  with initial segments  $\{x | x < a\}$  as elements of  $V_a$ . Consequently, on each initial segment, there are at most  $\aleph_1$  many possibilities for the sets coded by elements of  $S$ .

Let  $\sigma$  be the conjunction of the following statements:

- (A1) Both  $U(x, y)$  or  $V(x, y)$  imply  $x \in L$ . Writing  $U_x = \{y | U(x, y)\}$  and  $V_x = \{y | V(x, y)\}$ , the sets  $L, S, U_x, V_x$  (for all  $x \in L$ ) are pairwise disjoint and their union is everything.
- (A2) All  $E_n$  define equivalence relations on every set  $U_x$  and  $V_x$  where on every  $U_x$  or  $V_x$ ,  $E_0$  has exactly two classes and every  $E_n$ -class is the union of exactly two  $E_{n+1}$ -classes. In addition,  $\bigwedge_{n < \omega} x E_n y$  implies  $x = y$ .
- (A3)  $<$  is a linear ordering of  $L$ . For  $x \in L$  we write  $L_{<x} = \{y \in L | y < x\}$  and  $L_{\leq x} = L_{<x} \cup \{x\}$ .
- (A4)  $F(s, t, x)$  implies  $s, t \in S$  and  $x \in L$ .  $F$  defines a symmetric function from  $S \times S$  to  $L$ .
- (A5)  $R \subset S \times L$ . For  $s \in S$  we write  $R_s = \{x \in L | R(s, x)\}$ . For any two distinct  $s, t \in S$ ,  $R_s$  and  $R_t$  are identical on  $L_{\leq F(s,t)}$  and disjoint on  $L \setminus L_{\leq F(s,t)}$ .

- (A6)  $G(x, y, z)$  implies  $x \in L$ ,  $y < x$  and  $z \in U_x$ . For every  $x \in L$ ,  $G(x, \cdot, \cdot)$  defines a bijective function  $G_x : L_{<x} \rightarrow U_x$  by  $G_x(y) = z$  if and only if  $G(x, y, z)$ .
- (A7)  $H(x, y, z)$  implies  $x \in L$ ,  $y \in S$  and  $z \in V_x$ . For every  $x \in L$ ,  $H(x, \cdot, \cdot)$  defines a surjective function  $H_x : S \rightarrow V_x$  by  $H_x(y) = z$  if and only if  $H(x, y, z)$ .  $H_x$  has the property that  $H_x(s) = H_x(t)$  if and only if  $F(s, t) \geq x$ .

It is easy to construct a model of  $\sigma$ , but  $\sigma$  is not a complete sentence. We verify that it satisfies the hypotheses of Theorem 13. The axioms are all at most  $\Pi_2$ -statements, so we have preservation under chains. Also, since the equivalence relations  $E_n$  are refining and  $\mathcal{L} \setminus \{E_n\}_{n < \omega}$  is finite,  $S_{\text{qf}}(\sigma)$  is countable.

Towards showing EPC, let  $M \models \sigma$  be countable,  $\bar{a} = (a_1, \dots, a_n) \in M$  and let  $p(\bar{x}), q(\bar{x}, y) \in S_{\text{qf}}(\sigma)$  with  $\bar{a} \models p$  and  $p \subset q$  (note that it suffices to consider a single variable  $y$  instead of an arbitrary tuple). We want to find some countable  $N \supset M$  and  $b \in N$  such that  $\bar{a}b \models q$ . There are several cases:

- Suppose  $S(y) \in q(\bar{x}, y)$ . We will add a new point  $y$  to  $S$  and define a set  $R_y$  respecting the requirements of  $q$  and the axioms of  $\sigma$ . The requirements can be  $R(y, x_i)$ ,  $\neg R(y, x_i)$  as well as  $F(y, x_j) = x_i$ ,  $F(y, x_j) \neq x_i$  and  $H_{x_i}(y) = x_j$ ,  $H_{x_i}(y) \neq x_j$ ,  $H_{x_i}(y)E_n x_j$ ,  $\neg H_{x_i}(y)E_n x_j$  for components  $x_i, x_j$  in  $\bar{x}$  and  $n < \omega$  ( $G$  does not matter here since it does not involve elements from  $S$ ; also note that conditions like  $F(y, x_j) > x_i$  translate to  $F(y, x_j) = x_k \wedge x_k > x_i$  since  $F$  is not a function but a relation symbol).

Consider the set of all elements  $z \in L$  occurring in  $\bar{x}$  such that one of the following holds:

- (i)  $q \vdash F(y, x_i) = z$  for some  $x_i$  in  $\bar{x}$  or
- (ii)  $q \vdash R(y, z)$  and there is some  $s \in S$  with  $M \models R(s, z)$  or
- (iii)  $q \vdash H_z(y) = x_i$  and  $M \models H_z(s) = x_i$  for some  $x_i$  in  $\bar{x}$  and  $s \in S$ .

Let  $A = \{a \in L \mid q \vdash R(y, a)\}$ . We now have two cases:

- There is no such  $z$ . Then, we choose any  $c \in L$  that is smaller than any element of  $\bar{x}$ , as well as an arbitrary element  $s \in S$ . We set  $R_y = A \cup (R_s \cap L_{\leq c})$  and naturally  $F(y, s) = c$  (note that  $R_y$  and every  $R_t$  ( $t \in S$ ) are disjoint above  $c$  since (ii) fails).
- There is such an element and let  $z$  be the maximal such. We set  $R_y = A \cup (R_{x_i} \cap L_{\leq z})$  if  $z$  satisfies (i) and  $R_y = A \cup (R_s \cap L_{\leq z})$  in the cases (ii) and (iii) (choose any such  $s$  arbitrarily). If we are in the case (ii) or (iii) and  $q$  implies  $F(y, s) \neq z$ , we also add a new element  $w$  to  $L$  which is greater than  $z$  and smaller than any element of  $\bar{x}$  that is larger than  $z$ , and we declare  $R(s, w)$ ,  $R(y, w)$ ,  $F(s, y) = w$ .

In either of the two cases, we will have to turn  $M$  with the additional  $y$  (and possibly  $w$ ) into a model of  $\sigma$ . We have to set the  $F$ - and  $H$ -relations which can be done straightforwardly (respecting possible requirements from  $q$  for  $H$ ; we may

have to add new points to sets  $V_a$  for  $a > z$ ). In case we added the point  $w$ , we also have to add new sets  $U_w, V_w$  as well as a new point to each  $U_a$  for  $a > w$ , and extend  $G$  accordingly.

- Now suppose  $L(y) \in q(\bar{x}, y)$ . Add a new element  $z$  to  $L$  for  $y$  in an arbitrary cut that complies with the conditions  $x_i < y$  or  $x_i > y$  contained in  $q$ . Add  $R(x_i, z)$  whenever demanded by  $q$  and for any other  $s \in S$  add  $R(s, z)$  if and only if  $R(t, z)$  and  $F(s, t) > z$  for some element  $t \in S$ . Finally, we have to add new sets  $U_z$  and  $V_z$  as well as a new point  $a$  to each  $U_w$  with  $w > z$  and declare  $G(w, z, a)$ . We may have to add a new point to sets  $V_w$  for  $w > z$  too.
- Should  $U_{x_i}(y)$  or  $V_{x_i}(y)$  belong to  $q$ , it is easy to see that there must already be some  $b \in M$  with  $\bar{a}b \models q$ .

Now we apply Theorem 13 to  $\sigma$ . Immediately we see that  $\sigma^*$  implies:

- The ordering on  $L$  is dense without endpoints.
- Every set  $R_s$  is dense (and thus unbounded) and co-dense in  $L$ .
- $s \neq t$  implies  $R_s \neq R_t$  ("R is extensional").

But we know more about the properties of  $\sigma^*$ . The countable model of  $\sigma^*$  is extendible, so there is an uncountable model. In addition, we have seen in the verification of EPC that we have a lot of freedom in adding new elements to countable models of  $\sigma$ , and thus to models of  $\sigma^*$ , so that we can conclude the existence of models of  $\sigma^*$  with

- $(L, <)$  isomorphic to a proper initial segment of  $\eta_1 \cdot \omega_2$ , where  $\eta_1$  is the saturated dense linear order without endpoints of size  $\aleph_1$  (we assume GCH).
- all  $(U_x, E_n)_{n < \omega}$  and  $(V_x, E_n)_{n < \omega}$  isomorphic to  $(2^\omega, F_n)$  where we define  $\xi F_n \rho$  if and only if  $\xi(k) = \rho(k)$  for all  $k \leq n$ .

Now we consider the class  $\mathbb{P}$  of all such models with the additional properties

- $(L, <)$  is an initial segment of  $(\eta_1 \cdot \omega_2, <)$
- $S$  is a subset of  $\omega_3$  of size  $\aleph_1$  (so all models in  $\mathbb{P}$  will have size  $\aleph_1$ ).
- The sets  $U_x$  and  $V_x$  ( $x \in L$ ) equal  $2^\omega \times \{(x, 0)\}$  and  $2^\omega \times \{(x, 1)\}$  respectively and the  $E_n$  defined on them are the natural ones (compare with  $F_n$  above).

We order the elements of  $\mathbb{P}$  by the superstructure relation  $\supset$ . Since  $\sigma^*$  is preserved under unions, the poset  $(\mathbb{P}, \supset)$  is  $\omega_2$ -closed (meaning every sequence of length less than  $\omega_2$  of elements of  $\mathbb{P}$  has a lower  $\supset$ -bound; clearly the union of the chain of models will do).

Now we show that  $(\mathbb{P}, \supset)$  has the  $\omega_3$ -cc. Take any  $X \subset \mathbb{P}$  of size  $\aleph_3$ . We shall find two elements of  $X$  which have a common extension. By the pigeonhole-principle and the delta-system-lemma, we may assume that

- the collection of the underlying sets (of the models in  $X$ ) form a delta-system.
- the  $L$ -part of all models in  $X$  is identical.
- the  $\mathcal{L}$ -structure of all models in  $X$  is identical on the root of the delta-system.
- the collection of sets  $R_s$  ( $s \in S$ ) is identical for all elements of  $X$ .

Two models  $M, N \in X$  may only differ on their  $S$ -part. We would like to make the union  $M \cup N$  into a model of  $\sigma$ . The problem is that if the models are not already identical, there will be  $x \in S^M, y \in S^N$  outside the root such that  $R_x = R_y$ , so  $F(x, y)$  cannot be defined in such a way that axiom (A5) holds. The solution is to end-extend  $L$  in order to make  $R_x$  and  $R_y$  disjoint on a final segment.

Suppose that in  $\eta_1 \cdot \omega_2$ ,  $L$  is an initial segment contained in  $\{x \mid x < a\}$ . Enumerate the elements of  $S^M \setminus S^N$  as  $(s_\alpha)_{\alpha < \mu}$  (for some  $\mu \leq \aleph_1$ ). Now inductively do the following: given  $\alpha < \omega_1$  there is a unique  $t \in S^N \setminus S^M$  such that  $R_{s_\alpha} = R_t$ . Set  $R(s_\alpha, a), R(t, a), F(s_\alpha, t) = a$  and  $R(s_\alpha, a_\alpha)$  (but *not*  $R(t, a_\alpha)$ ), where  $a_\alpha \in \eta_1 \cdot \omega_2$  is greater than  $a$  and any already chosen  $a_\beta$  ( $\beta < \alpha$ ). Now we have to add sets  $U_{a_\alpha}$  and  $V_{a_\alpha}$  and extend  $G$  and  $H$  to get a model  $M'$  of  $\sigma$  containing both  $M$  and  $N$ . We do so in a way that it is still possible to add  $\aleph_1$  more points to those sets  $U_{a_\alpha}$  and  $V_{a_\alpha}$  later on (which will be necessary when we construct an extension satisfying  $\sigma^*$ , see below). Note that we do not have to add any point to the  $U_x, V_x$  for  $x \in L^M$  which is fortunate since that would be impossible.

Having obtained a model  $M'$  of  $\sigma$  containing both  $M$  and  $N$  as submodels, our final task in proving  $\omega_3$ -cc is to extend  $M'$  to an element  $M''$  of  $\mathbb{P}$ . In particular, we want  $M''$  to have the following properties:

- $M''$  must be a model of  $\sigma^*$ .
- $L^{M''}$  must be an initial segment of  $\eta_1 \cdot \omega_2$ .
- The sets  $U_x^{M''}$  and  $V_x^{M''}$  must be equal to  $2^\omega \times \{(x, 0)\}$  and  $2^\omega \times \{(x, 1)\}$  respectively.

We will construct a continuous chain  $(M_\alpha)_{\alpha < \omega_1}$  of models of  $\sigma$  starting from  $M_0 = M'$  such that  $M'' = \bigcup_{\alpha < \omega_1} M_\alpha$  satisfies our requirements. We have several sets of “tasks”

(each enumerated in ordertype  $\omega_1$ ) that we want to perform along that chain:

- Let  $W$  be an enumerated set of the tasks “add the element  $w$  to the  $L$ -part of the so far constructed model” for any  $w \in \eta_1 \cdot \omega_2$  that is smaller than some  $a_\alpha$  (we constructed the elements  $a_\alpha$  above). Thus after performing all tasks in  $W$ , the  $L$ -part of  $M''$  will be an initial segment of  $\eta_1 \cdot \omega_2$  (the smallest one containing all  $a_\alpha$ ).
- Having reached stage  $\alpha$  of the chain, enumerate all pairs  $(\bar{a}, q)$  with  $\bar{a} \in M_\alpha$  and  $\text{tp}_{\text{qf}}(\bar{a}) \subset q \in S_{\text{qf}}(\sigma)$  as  $T_\alpha = ((\bar{a}_\beta, q_\beta))_{\beta < \omega_1}$  (compare with the proof of Theorem 13). Designate the set of tasks “add a tuple  $\bar{b}$  such that  $\bar{a}_\beta \bar{b} \models q_\beta$ ” by  $T_\alpha$ .

- Having reached stage  $\alpha$  of the chain, let  $X_\alpha$  and  $Y_\alpha$  respectively be enumerated sets of the tasks “add the element  $(\rho, (x, 0))$  to  $U_x$ ” and “add the element  $(\rho, (x, 1))$  to  $V_x$ ” for all  $\rho \in 2^\omega$  and  $x \in L^{M_\alpha}$ .

At each stage  $\alpha$  of the chain, add elements to the model  $M_\alpha$  such that the least task in  $W$  as well as in all  $T_\beta, X_\beta, Y_\beta$  ( $\beta \leq \alpha$ ) is performed. Then remove those tasks from the sets  $W, T_\beta, X_\beta, Y_\beta$ . By possibly adding additional elements, we can do this while obtaining a model  $M_{\alpha+1}$  of  $\sigma$  (the arguments and techniques are the same as in the proof that  $\sigma$  has EPC). Again, when we add new sets  $U_w$  and  $V_w$ , we do so in a way that it is still possible to add  $\aleph_1$  many additional elements to them later on.

Note that at each stage, we only have to add countably many elements in each  $U_x, V_x$  and the  $L$ -part of  $M_\alpha$ , so we do not encounter the problem of saturating at a countable stage of the chain-construction the  $U_x, V_x$  for  $x > a$  ( $a$  as defined above) or any part of  $L$  above  $a$ . This would be a serious problem as for example adding a new element  $w$  to the order requires adding a *new* element to the  $U_x$  with  $x > w$  (because of the properties of  $G$ ). We thus are able to carry out the construction through all countable ordinals and obtain  $M''$  as the union of the chain with the required properties. This concludes the proof of  $\omega_3$ -cc.

Let  $G$  be a  $\mathbb{P}$ -generic filter over  $V$ .  $\bigcup G$  will be a model of  $\sigma^*$  of size  $\aleph_3^V$ . But since the forcing is  $\omega_2$ -closed and has  $\omega_3$ -cc, all cardinals are preserved and in particular  $\aleph_3^{V[G]} = \aleph_3^V$ . I.e. we get a model of  $\sigma^*$  of size  $\aleph_3$  in a generic extension. On the other hand, any such model codes an  $\aleph_2$  Kurepa family which means that it is consistent with ZFC+GCH (assuming the existence of an inaccessible cardinal and noting that the forcing preserves GCH<sup>2</sup>) that  $\sigma^*$  has no model of size  $\aleph_3$ .

## 5 Final observations

The question of absoluteness of model-existence (under ZFC+GCH) in  $\aleph_\omega$  remains open. On the other hand, the technique of finding complete examples described in section 4 should be applicable more widely to obtain complete examples of non-absoluteness of model existence (under ZFC+GCH) in cardinals greater than  $\aleph_3$ . Interestingly, however, this method seems to be problematic for finding examples for model existence in  $\aleph_2$ , at

---

<sup>2</sup>This is a standard argument: For any (infinite)  $\kappa$ , each subset of  $\kappa$  added by the forcing is of the form

$$\{\alpha < \kappa \mid G \cap A_\alpha \text{ is nonempty}\}$$

where  $\vec{A} = (A_\alpha)_{\alpha < \kappa}$  is in the ground model and each  $A_\alpha$  is an antichain in the forcing. (This is because we can take a name  $\sigma$  for the given set, let  $B_\alpha$  be a maximal antichain consisting of conditions which decide “ $\alpha \in \sigma$ ” and take  $A_\alpha$  to consist of the elements of  $B_\alpha$  which force “ $\alpha \in \sigma$ ”.)

As GCH holds in the ground model and the forcing has  $\omega_3$ -cc, the fact that the forcing has size  $\omega_3$  implies that there are only  $((\omega_3)^{\omega_2})^\kappa = (\omega_3)^\kappa$  many (in the sense of the ground model) such sequences  $\vec{A}$ . For  $\kappa \geq \omega_2$ , this is  $2^\kappa = \kappa^+$ . As the forcing does not add subsets of  $\omega_1$ , the GCH will also hold at  $\omega_1$ .

least with the approach of trying to code Kurepa families. The reason is that it seems difficult to code an  $\omega_1$ -like ordering without making many elements definable over others (or even getting infinite definable closures over finite tuples), which destroys any chance to have EPC.

As a last remark, our use of the concept of Kurepa families has the slight flaw that in order to find set-theoretic universes which do not contain such families, we have to assume the existence of inaccessible cardinals. For the special Aronszajn technique, we even have to assume the consistency of supercompact cardinals. It would be nice to find  $L_{\omega_1, \omega}$  sentences for which under GCH the existence of models of certain cardinalities is not absolute, without assuming the existence of large cardinals.

### Acknowledgements

We would like to express our gratitude to the John Templeton Foundation for supporting the Infinity Project (hosted by the CRM in Barcelona) in which the presented work has been accomplished. We also greatly thank John Baldwin, Fred Drucek, Rami Grossberg and Andrés Villaveces for helpful discussions at the CRM, especially on the idea of coding Kurepa families into models to show non-absoluteness of model existence. Rami Grossberg in particular suggested generalizing a theory coding an  $\aleph_1$  Kurepa family to higher cardinalities to achieve non-absoluteness of model existence in cardinalities above  $\aleph_2$ . He also was the first to recognize a problem in dealing with  $\aleph_\omega$  by this construction.

Sy-David Friedman  
Kurt Gödel Research Center for Mathematical Logic  
Währinger Strasse 25, 1090 Wien, Austria  
sdf@logic.univie.ac.at

Tapani Hyttinen  
Department of Mathematics University of Helsinki  
P.O. Box 68 (Gustaf Hällströmin katu 2b), FI-00014 Helsinki  
tapani.hyttinen@helsinki.fi

Martin Koerwien  
Kurt Gödel Research Center for Mathematical Logic  
Währinger Strasse 25, 1090 Wien, Austria  
koerwien@math.uic.edu

### References

- [1] Baldwin, J. T., *Amalgamation, absoluteness and categoricity*, Proceedings of Southeast Asia Logic Conference (June 2009), to appear. Available at

www.math.uic.edu/jbaldwin.

- [2] Barwise, K. J., *The role of the omitting types theorem in infinitary logic*, Archiv für Mathematische Logik und Grundlagenforschung, vol. 21, 1981, pp. 55-68.
- [3] Chang, C. C. and Keisler, H. J., *Model Theory*, North Holland, 1990.
- [4] Cummings, J.; Foreman, M.; Magidor, M. *Squares, scales and stationary reflection*, Journal of Mathematical Logic, vol. 1, 2001, pp. 35-98.
- [5] Gao, S. *On automorphism groups of countable structures*, The Journal of Symbolic Logic, vol. 63(3), 1998, pp. 891-896.
- [6] Gregory, J., *Elementary extensions and uncountable models for infinitary finite quantifier language fragments. Preliminary report*, Notices of the American Mathematical Society, vol. 17, 1970, pp. 967,968.
- [7] Grossberg, R., Shelah, S. *On the number of non isomorphic models of an infinitary theory which has the order property part A.*, Journal of Symbolic Logic, vol. 51, 1986, pp. 302-322.
- [8] Hjorth, G. *Knight's model, its automorphism group, and characterizing the uncountable cardinals*, Journal of Mathematical Logic, vol. 2(1), 2002, pp. 113-144.
- [9] Jech, T., *Set Theory*, Springer Monographs in Mathematics, 2002.
- [10] Jensen, R. B., *The fine structure of the constructible hierarchy. With a section by Jack Silver*, Annals of Mathematical Logic, vol. 4, 1972, pp. 229-308.
- [11] Keisler, H. J., *Model Theory for Infinitary Logic*, North-Holland Publishing Company, 1971.
- [12] Malitz, J., *The Hanf number for complete  $L_{\omega_1, \omega}$ -sentences*, in The syntax and semantics of infinitary languages (J. Barwise, Editor), Lecture Notes in Mathematics, vol. 72, Springer Edition, 1968, pp. 166-181.
- [13] Specker, E. *Sur un problème de Sikorski*, Colloq. Math. 2, 1949, pp. 9-12.